

**Program in Applied Mathematics**  
**PROBABILITY AND STATISTICS PRELIMINARY EXAMINATION**  
**January 2009**

<b>Notice.</b> Do four of the following five problems. Place an X on the line opposite the number of the problem that you are <b>NOT</b> submitting for grading. Please do not write your name anywhere on this exam. You will be identified only by your student number, given below and on each page submitted for grading. Show all relevant work!	1. ___ 2. ___ 3. ___ 4. ___ 5. ___ <b>TOTAL.</b> ___
--	---

**STUDENT NUMBER:** \_\_\_\_\_

1. Imagine rolling an unbiased die repeatedly and let  $T_k$  be the number of rolls until some number appears repeated  $k$  times in a row. For example, if the first few rolls of the die were 6;4;5;5;5;4;... then  $T_1 = 1$ ,  $T_2 = 4$ , and  $T_3 = 5$ . (Note that  $T_1$  is always 1.) Find a formula that relates  $E(T_k)$  to  $E(T_{k-1})$  and use it to determine  $E(T_k)$  explicitly for all  $k \geq 1$ .
2. This problem has two separate parts.
  - (a) (*Markov's inequality.*) Show that  $a^2 \cdot P[|X| \geq a] \leq E(X^2)$ , for any random variable  $X$  and real-constant  $a > 0$ .
  - (b) Consider random variables  $T_1; T_2; T_3; \dots$  i.i.d. Exponential(1) i.e. each has probability density function  $f(t) = e^{-t}$ , for  $t \geq 0$ . Consider a certain continuous function  $g: \mathbb{R} \rightarrow \mathbb{R}$

(c) Motivated by part (b) consider the random variable

$$Z := \begin{cases} Y = (1 - \alpha)^{1-n} & ; Y \leq (1 - \alpha)^{1-n}; \\ Y & ; Y > (1 - \alpha)^{1-n}; \end{cases}$$

Show that  $Z$  is a lower-bound for  $\alpha$  with a confidence of at least  $100(1 - \alpha)\%$ .

4. Let  $0 < p < 1$  be an unknown parameter and consider a random sample  $(X_1; X_2)$  such that  $P[X_1 = k_1; X_2 = k_2] = p^2 \cdot (1 - p)^{k_1 + k_2}$ , for  $k_1; k_2 \geq 0$  integers. In what follows we will say that a real-valued function  $g(p)$  is *good* if there are constants  $c_0; c_1; c_2; \dots$  such that  $g(p) = \sum_{k=0}^{\infty} c_k \cdot (1 - p)^k$ , for all  $0 < p < 1$ .
  - (a) Show that if  $g(p)$  is a good function and  $c_0; c_1; c_2; \dots$  are like above then  $X_1$  is an unbiased statistic for  $g(p)$ .
  - (b) Find a UMVUE based on  $(X_1; X_2)$  for any good function  $g(p)$ .
  - (c) Find the UMVUE based on  $(X_1; X_2)$  for  $g(p) = p(1 + p)$ .
  
5. Consider a fleet of  $N$  buses each of which breaks down independently of the others at a rate  $\lambda$ . When a bus brakes down, it is sent for repair to a depot. The mechanic of the depot can only repair one bus at a time and the repair time is always an exponential random variable with mean  $1 = \mu$ .
  - (a) Determine the equilibrium distribution of the number of buses in the repair depot (i.e. undergoing repair or waiting to be repaired).
  - (b) If at a given moment all buses are functional, what is the probability that in the next  $t$  units of time no bus will break down? Determine this probability explicitly.
  - (c) Assume that  $N = 1$ . If at a given moment all buses are functional, what is the probability that  $t$  units of time later no bus will be in the repair depot? Determine this probability explicitly.